

From Reissner-Nordström quantum states to charged black holes mass evaporation

P.V. Moniz

*DAMTP, University of Cambridge
Silver Street, Cambridge, CB3 9EW, UK*

e-mail: prlv10@amtp.cam.ac.uk; paul.vargasmoniz@ukonline.co.uk

WWW-site: <http://www.damtp.cam.ac.uk/user/prlv10>

In this report we describe quantum Reissner-Nordström (RN) black-holes interacting with a complex scalar field. Our analysis is characterized by solving a Wheeler-DeWitt equation in the proximity of an apparent horizon of the RN space-time. Subsequently, we obtain a wave-function $\Psi_{\text{RN}}[M, Q]$ representing the RN black-hole. A special emphasis is given to the evolution of the mass-charge rate affected by Hawking radiation. More details can be found in ref. ¹².

Recently, there has been a renewed interest in the canonical quantization of black-hole space-times ¹⁻¹². The general aim is to obtain a description of quantum black holes that would go beyond a semi-classical approximation. In this report, we will extend M. Pollock's method ¹ (which was itself influenced by the work of A. Tomimatsu ⁴) to Reissner-Nordström (RN) black-hole. As a consequence, we will find a wave function for the RN black-hole, which will have an explicit dependence on its mass M and the charge Q . The T.O.H. (Tomimatsu-Oda-Hosoya) method ^{5,6} constitutes another very interesting and similar approach to this purpose.

The main elements of the Hamiltonian formalism used here follow ref. ¹³. En route towards our reduced model we further take the following steps. To begin with, our 4-dimensional *spherically symmetric* metric is written as

$$ds^2 = h_{ab}dx^a dx^b + \phi^2(d\theta^2 + \sin^2\theta d\varphi^2), \quad (1)$$

together with the ADM decomposition

$$h_{ab} = \begin{pmatrix} -\alpha^2 + \frac{\beta^2}{\gamma} & \beta \\ \beta & \gamma \end{pmatrix}, \quad h^{ab} = \begin{pmatrix} -\frac{1}{\alpha^2} & \frac{\beta}{\alpha^2\gamma} \\ \frac{\beta}{\alpha^2\gamma} & \frac{1}{\alpha} - \frac{\beta^2}{\alpha^2\gamma^2} \end{pmatrix}, \quad (2)$$

where $\alpha, \beta, \gamma, \phi$ are functions of $(x^0, x^1) = (\tau, r)$ which will be defined later (see eq. (3)). We also take a scalar field $\hat{\psi} = \psi(\tau, r)$ and the electromagnetic field $F_{ab} = \varepsilon_{ab}\sqrt{-h}E$; $E = (-h)^{-1/2}(\dot{A}_1 - A'_0)$, $\hat{D}_a\hat{\psi} = \partial_a\psi + ieA_a\psi$; $\hat{D}_2\hat{\psi} = \hat{D}_3\hat{\psi} = 0$. Notice we employ “.” $\equiv \frac{\partial}{\partial\tau}$ and “'” $\equiv \frac{\partial}{\partial r}$. The next *significant* step consists in choosing the following coordinate gauge: $\alpha = \frac{1}{\sqrt{\gamma}} \Leftrightarrow \sqrt{-h} = 1$.

The constraints of our model will be expressed in terms of dynamical quantities defined at an apparent horizon of the RN black-hole, which is defined by the condition¹³: $h^{ab}(\partial_a\phi)(\partial_b\phi) = 0 \Leftrightarrow \dot{\phi}(\dot{\phi} - \phi') = 0$. In order to obtain a satisfactory description of an evaporating RN black hole on the apparent horizon, it is more convenient to use a RN - Vaidya metric ¹. With $v \equiv \tau + r = t + r^*$ as the advanced

null Eddington-Finkelstein coordinate, and r^* as the corresponding “tortoise” coordinate, the relationship between the time τ and the time coordinate t for the standard RN metric is $\tau = t - r + r^*$. The Vaidya RN metric becomes

$$ds^2 = - \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right) d\tau^2 + \left(\frac{4M}{r} - \frac{2Q^2}{r^2}\right) d\tau dr + \left(1 + \frac{2M}{r} - \frac{Q^2}{r^2}\right) dr^2 + \phi^2 d\Omega_2^2. \quad (3)$$

The RN black-hole has two apparent horizons, namely at $r_{\pm}(v) = M(v) \pm \sqrt{M^2(v) - Q^2}$ and we will henceforth restrict ourselves to the case of r_+ . In similarity with the Schwarzschild case we also take $M \simeq M(\tau)$ in the vicinity of the apparent horizon. After some lengthy calculations we obtain (see ref. ¹² for more details) the approximate expressions:

$$\begin{aligned} \mathcal{H}_0 &= \frac{1}{2}\pi_\phi - \frac{1}{8} + \frac{Q^2}{32M^2} - \frac{1}{16\pi}\frac{1}{\rho^2}(\pi_\chi^2 + \pi_{\psi_1}^2) - \rho^2\pi(\psi_1'^2 + \chi'^2) \\ &\quad - \frac{1}{2\rho^2}\pi_{A_1}^2 - \frac{\rho^2}{2}[2\pi e^2 A_1^2(\psi_1^2 + \chi^2)] - 2\pi e\rho^2 A_1(\psi_1\chi' - \chi\psi_1') \end{aligned} \quad (4)$$

$$\mathcal{H}_1 = \frac{1}{2}\pi_\phi - \frac{1}{8} - \frac{3Q^2}{32M^2} + \frac{1}{2}[\pi_{\psi_1}\psi_1' + \pi_\chi\chi'] + \frac{eA_1}{4}(\pi_{\psi_1}\chi - \pi_\chi\psi_1), \quad (5)$$

with $\psi = \sqrt{2\pi}(\psi_1 + i\chi)$, $\pi_\psi \rightarrow \frac{1}{\sqrt{8\pi}}\pi_\psi$. Compatibility¹² requires that $\pi_{\psi_1} = -4\pi\rho^2\psi_1'$, $\pi_\chi = -4\pi\rho^2\chi'$, with the terms with A_1 to be negligible, and with the conditions $A_0 = 0$ together with $\dot{\psi} = \dot{\psi}^\dagger = 0$. We further take $A_1 = A_1(r)$.

Quantization proceeds via the operator replacements

$$\pi_\phi \rightarrow -i\frac{\partial}{\partial\phi} \simeq -i\frac{2M^2}{4M^2 + Q^2}\frac{\partial}{\partial M}, \pi_{\psi_1} \rightarrow -i\frac{\partial}{\partial\psi_1}, \pi_\chi \rightarrow -i\frac{\partial}{\partial\chi}, \quad (6)$$

which yields the Wheeler-DeWitt equation for the wave function Ψ ,

$$-i\frac{2M^2}{4M^2 + Q^2}\frac{\partial\Psi}{\partial M} = -\frac{1}{16\pi M^2 + \pi Q^4/M^2 - 8\pi Q^2}\left[\frac{\partial^2\Psi}{\partial\psi_1^2} + \frac{\partial^2\Psi}{\partial\chi^2}\right] + \frac{1}{4}\Psi, \quad (7)$$

whose solutions are

$$\Psi_{\text{RN}}[M, Q; \psi_1, \chi; k] = \Psi_{\text{RN}}^0 e^{i\left[\frac{1}{4}\left(-\frac{Q}{M} + 2M\right) + k^2\frac{M}{Q^2 - M^2} \pm 2\sqrt{\pi}k(\psi_1 + \chi)\right]} \zeta(A_1), \quad (8)$$

where k^2 is a separation constant and Ψ_{RN}^0 an integration constant.

It is interesting to notice the following as well. For the Schwarzschild case ($Q = 0$) eq. (8) implies that near to $M = 0$ the wave function will oscillate with infinite frequency. If $\dot{M} < 0$, this would represent the quantum mechanical behaviour of the black hole near the end point of its evaporation. In the RN case, the rapid oscillations will occur again for $M = 0$ but also when $M \sim Q$. I.e., near extremality and when the black hole mass evaporation can eventually stop. Hence, the presence of Q in Ψ_{RN} allowed us to identify some known physical situations of the RN black hole.

As far as the mass-charge ratio for the RN black hole is concerned, we get the equation

$$\dot{M} + \frac{1}{4M^2} \frac{a[k^2; Q]}{d(M)} + \frac{1}{4M^4} \frac{b[k^2; Q]}{d(M)} + \frac{c[k^2; Q]}{d(M)M^6} = 0, \quad (9)$$

where

$$a[k^2; Q] = k^2 - 5Q^2 + Q; b[k^2; Q] = k^2Q^2 - 3Q^4; c[k^2; Q] = \frac{k^2Q^4}{8}; d(M) = 1 + \frac{Q^2}{2M^2}. \quad (10)$$

An integration of (9) leads to the result

$$M = \left[M_0^3 - \frac{3}{2}(k^2 - 5Q^2 + Q) \right]^{1/3} (t - t_0)^{1/3}. \quad (11)$$

We can now identify several physical cases of interest for the RN black hole, according if a, b, c, d are either positive, zero or negative.

For the case of $Q = 0$ ¹ (Schwarzschild), it is the separation constant k^2 that determines if the black hole is evaporating and decreasing its mass ($k^2 > 0$), or increasing its mass ($k^2 < 0 \Leftrightarrow k$ imaginary). In the present RN black hole case, the presence of the charge Q introduces significant changes. A $\dot{M} > 0$ stage can be obtained, with $k^2 > 0$ but $a < 0, b < 0$ and $c > 0, d > 0$. If $Q = 0$, this possibility is absent. When $Q \neq 0, k^2 < 0$, then $c < 0, b < 0, a < 0$, if $1 + 20k^2 > 0$ has real solutions.

Overall, our results do bring additional information regarding quantum black-hole, but there is still the need for further investigation.

Acknowledgments

This work was supported by the JNICT/PRAXIS XXI Fellowship BPD/6095/95. The author is grateful to M. Cavàglia, A. Hosoya, J. Louko and M. Ryan for interesting conversations and discussions, as well as to J. Mäkelä for further important comments and suggestions. He also wishes to thank I. Oda for letting him know of his recent work and for extensive correspondence.

References

1. M. Pollock, Int. J. Mod. Phys. **D3** (1994) 579
2. M. Cavaglia, V. De Alfaro and A. Filipov, Int. J. Mod. Phys. **D4** (1995) 661
3. M. Ryan and O. Obregon, talk given at the MG8 conference.
4. A. Tomimatsu, Phys. Lett. **B289** (1992) 283
5. A. Hosoya and I. Oda, gr-qc/9610055, submitted to Prog. Theo. Physics.
6. I. Oda, gr-qc/9703056, 9701058, 9704021, 9703055
7. K. Kuchar, Phys. **D50** (1994) 3961
8. J. Louko and S. Winters-Hilt, Phys. Rev. **D54** (1996) 2647
9. J. Mäkelä, Phys. Lett. **B390** (1996) 115
10. J. Louko and J. Mäkelä, Phys. Rev. **D54** (1996) 4982
11. C. Kiefer and T. Brotz, Phys. Rev. **D55** (1997) 2186

12. P. V. Moniz, *Wave function for the Reissner-Nordstrom black hole*, Mod. Phys. Lett. **A** (1997), to appear.
13. P. Thoni et al, Phys. Rev. **D30** (1984) 1168, P. Hajicek, Phys. Rev. **D30** (1984) 1178